

Brown, Culum (2016) Fish pain: An inconvenient truth. Animal Sentience 3(32) DOI: 10.51291/2377-7478.1069 Date of submission: 2015-11-30 Date of acceptance: 2015-12-13



This article has appeared in the journal *Animal Sentience*, a peer-reviewed journal on animal cognition and feeling. It has been made open access, free for all, by WellBeing International and deposited in the WBI Studies Repository. For more information, please contact wbisr-info@wellbeingintl.org.



Fish pain: An inconvenient truth

Commentary II on Key on Fish Pain

Culum Brown Biological Sciences Macquarie University

Abstract: Whether fish feel pain is a hot political topic. The consequences of our denial are huge given the billions of fish that are slaughtered annually for human consumption. The economic costs of changing our commercial fishery harvest practices are also likely to be great. Key outlines a structure-function analogy of pain in humans, tries to force that template on the rest of the vertebrate kingdom, and fails. His target article has so far elicited 34 commentaries from scientific experts from a broad range of disciplines; only three of these support his position. The broad consensus from the scientific community is that fish most likely feel pain and it is time governments display courage enough to act.

Culum Brown <u>CulumBrown@yahoo.com</u> studies the behavioural ecology of fishes with a special interest in learning and memory. He is Associate Professor of vertebrate evolution at Macquarie University, Co-Editor of the volume *Fish Cognition and Behavior*, and Editor for *Animal Behaviour* of the *Journal of Fish Biology*. <u>https://sites.google.com/site/culumbrown/</u>



Key (Key 2016; Key 2016a; Key 2016b) has put forth a structure-function analogy of pain in humans and then sets out to discover how far one might take that template when applied to the rest of the vertebrate kingdom – well, fish and the odd rodent at any rate. More than 30 commenters responded to the article, and this clearly shows that this topic is still controversial. Of these, three (Rose; Hart; Diggles) support Key's position. The vast majority of commentaries, however, do not, and argue that fish most likely feel pain. Most agree that Key's argument is flawed at best, and his evidence of how pain works in humans is selective, simplistic, misleading and outdated (Damasio & Damasio; Merker; Panksepp; Shriver). One emerging consensus, however, is that no single line of scientific evidence should over-rule any other, and many of the respondents agree that behavioural studies, such as conditioned place aversion, are an important component to understanding pain in human and non-human vertebrates alike.

The primary message from these commentaries is that Key's argument is fundamentally flawed from an evolutionary perspective. He argues (although later denies it) that human brain architecture is required to feel pain. The mechanistic approach centres around the role of the cortex in human pain. But as <u>Dinets</u>, <u>Brown</u>, <u>Ng</u> and others point out, the human cortex has taken on a huge number of roles that once were the domain of other brain regions. To suggest that fish don't feel pain because they lack a cortex, one would also have to write-off consciousness (<u>Seth</u>) or indeed any cognitive function that occurs in the human

cortex. One example of applying this logic would be to conclude that fish are not capable of learning either. Clearly this argument is absurd (see Brown et al. 2011; Brown 2015 for reviews). Jones, Mather, Striedter, Elwood and Edelman all point out that Key's approach also denies any possibility of convergent evolution, but surprisingly Key accepts that birds might provide an example of an independent evolution of a cortex-like structure. Haikonen, Manzotti, and Seth take it one step further and suggest that we don't even understand pain in humans yet, and it is far too early to be making judgement calls on other taxa. The role of the cortex in human pain is still debated (Stevens; Damasio & Damasio) and is most likely not the "on-off switch" for consciousness (Segner). Walters argues that high levels of consciousness are not required for pain perception, a view which is consistent with evolution being a gradual process rather than occurring as all-or-nothing leaps and with the notion that pain is likely an ancient evolutionary trait.

Broom, **Sneddon & Leach**, and Brown (2015) stated that the separation of nociception from the emotional response to it (pain) is old-fashioned and somewhat counter-productive. They argue that nociception and pain are part and parcel of the same system, and given the important nature of its function (harm avoidance), this system is likely evolutionarily ancient. It should hardly be surprising if this system is highly conserved across all vertebrates given the associated fitness advantages (<u>Seth</u>; <u>Striedter</u>; <u>Elwood</u>). The fact that the brain regions that are responsible vary across taxa is consistent with what we know to be the case with many other brain functions.

Diggles claims that the cheerleaders of fish pain perception (and welfare generally) intrinsically link pain with welfare. He rightly points out that other measures of welfare are available and are far less controversial. Of course the reason we are not talking about these other measures here is because they have been the foundation of fish welfare for decades (see Huntingford et al. 2006). There is no argument to be had there. The question we are addressing here is whether fish feel pain and whether we should include it in our welfare framework as we do with other vertebrates. Everyone agrees that current commercial harvesting methods are highly stressful for fish. Sadly, we currently lack the political will to do anything about it. I hope that given the known relationship between stress levels and filet quality, the aquaculture industry will move to limit pain and stress in fishes in the absence of any legislation (i.e., self-regulation driven by economics). Unfortunately, **Diggles** also needs a refresher course in vertebrate evolution since he wrongly states that fish are more closely related to sharks than they are to mammals and uses that as a basis for likening pain in fish to sharks rather than to mammals. It is true that "human-like" nociceptors have not been identified in elasmobranchs yet, but that is not to say that they do not exist. Science 101: Absence of evidence is not evidence of absence.

<u>Hart</u> rightly points out that there are issues with terminology and that humans are selfobsessed. Frankly, I think the whole consciousness debate is distracting and is not likely to be solved any time soon. We still don't really know what it is or how it is generated in humans let alone in nonhuman animals. Nevertheless, recent papers (e.g., Rey et al. 2015) certainly suggest that fish are conscious beings based on any criteria currently in use. <u>Wadiwel</u> suggests we should just accept that we cannot experience what other animals (or people) are feeling and that the question of pain in fish is hence one of making cautious ethical decisions under conditions of uncertainty: the weight of the evidence for and against pain in fish, and the weight of the consequences of a false negative (inferring fish do not feel pain when in reality they do).

The vast majority of commentaries – experts in a wide spectrum of relevant scientific and ethical specialities – accept the accumulating, multi-disciplinary evidence that it is likely that fish feel pain; moreover, even in the case of the fence-sitters (including optimistic agnostics), it is quite apparent that the precautionary principles apply, given the monumental number of fish killed each year in commercial fisheries. Indeed, <u>Jones</u> suggests that such a position is not only prudent, but also ethically obligatory. We should not allow conclusions drawn from scientific evidence to be tainted by the potential commercial consequences, as Key seems to recommend.

References

- Broom, D.M. (2016). Fish brains and behaviour indicate capacity for feeling pain. Animal Sentience 2016.010
- Brown, C. (2015). Fish intelligence, sentience and ethics. Animal Cognition, 18, 1–17
- Brown, C. (2016). Comparative evolutionary approach to pain perception in fishes. *Animal Sentience* 2016.011
- Brown, C. Krause, J. & Laland, K. (2011). *Fish Cognition and Behavior* (2nd Edition). Oxford: Wiley-Blackwell.
- Damasio, A. & Damasio, H. (2016). Pain and other feelings in humans and animals. *Animal Sentience* 2016.059
- Diggles, B.K. (2016). Fish pain: Would it change current best practice in the real world? Animal Sentience 2016.061
- Dinets, V. (2016). No cortex no cry. Animal Sentience 2016.013
- Edelman, D.B. (2016). Leaving the door open for fish pain: evolutionary convergence and the utility of 'just-so stories.' *Animal Sentience* 2016.062
- <u>Elwood, R.W. (2016)</u>. A single strand of argument with unfounded conclusion. *Animal* Sentience 2016.026
- Haikonen, P.O. (2016). On the sentience of fish. Animal Sentience 2016.014
- Hart, P.J.B. (2016). Fighting forms of expression. Animal Sentience 2016.015
- Huntingford, F. A., Adams, C., Braithwaite, V. A., Kadri, S., Pottinger, T. G., Sandøe, P. & Turnbull, J. F. (2006). Current issues in fish welfare. *Journal of Fish Biology*, 68, 332–372.

- Jones, R.C. (2016). Fish sentience and the precautionary principle. *Animal Sentience* 2016.016
- Key, B. (2016). Why fish do not feel pain. Animal Sentience 2016.003
- Key, B. (2016a). Going beyond just-so stories. Animal Sentience 2016.022
- Key, B. (2016b). Falsifying the null hypothesis that "fish do not feel pain." Animal Sentience 2016.039
- Manzotti, R. (2016). No evidence that pain is painful neural process. *Animal Sentience* 2016.017
- <u>Mather, J.A. (2016)</u>. A wider view of the pain problem from an invertebrate perspective. *Animal Sentience* 2016.018
- Merker, B. (2016). Drawing the line on pain. Animal Sentience 2016.030
- Ng, Y.-K. (2016). Could fish feel pain? A case for a wider, multi-faceted perspective. Animal Sentience 2016.019
- Panksepp, J. (2016). Brain processes for "good" and "bad" feelings: how far back in evolution? *Animal Sentience* 2016.031
- Rey, S., Huntingford, F.A., Boltaña, S., Vargas, S., Knowles, T.G. & Mackenzie, S. (2015). Fish can show emotional fever: stress-induced hyperthermia in zebrafish. *Proceedings of the Royal Society of London B: Biological Sciences*, 282(1819).
- Rose, J. D. (2016). Weighing the evidence. Animal Sentience 2016.032.
- <u>Segner, H. (2016)</u>. Why babies do not feel pain, or: how structure-derived functional intepretations can go wrong. *Animal Sentience* 2016.033
- <u>Seth</u>, A.K. (2016). Why fish pain cannot and should not be ruled out. *Animal Sentience* 2016.020
- <u>Shriver, A. (2016)</u>. Cortex necessary for pain not in the sense that it matters. *Animal* Sentience 2016.034
- Sneddon, L.U. & Leach, M.C. (2016). Anthropomorphic denial of fish pain. Animal Sentience 2016.035
- Stevens, E.D. (2016). Why is fish "feeling" pain controversial? Animal Sentience 2016.036
- <u>Striedter, G. (2016)</u>. Lack of neocortex does not imply fish cannot feel pain. *Animal Sentience* 2016.021

Wadiwel, D. (2016). Fish and pain: the politics of doubt. Animal Sentience 2016.038

Walters, E.T. (2016). Pain-capable neural substrates may be widely available in the animal kingdom. *Animal Sentience* 2016.063